

Dynamics in a supercooled liquid of symmetric dumbbells: Reorientational hopping for small molecular elongations

Angel J. Moreno,^{1,2} Song-Ho Chong,³ Walter Kob,⁴ and Francesco Sciortino¹

¹*Dipartimento di Fisica and INFM-CRS-SOFT, Università di Roma La Sapienza, P.le. A. Moro 2, 00185 Roma, Italy*

²*Donostia International Physics Center, Paseo Manuel de Lardizabal 4, 20018 San Sebastián, Spain*

³*Institute for Molecular Science, Okazaki 444-8585, Japan*

⁴*Laboratoire des Colloïdes, Verres et Nanomatériaux,*

Université Montpellier II and UMR 5587 CNRS, 34095 Montpellier Cedex 5, France

We present extensive molecular dynamics simulations of a liquid of symmetric dumbbells, for constant packing fraction, as a function of temperature and molecular elongation. For large elongations, translational and rotational degrees of freedom freeze at the same temperature. For small elongations only the even rotational degrees of freedom remain coupled to translational motions and arrest at a finite common temperature. The odd rotational degrees of freedom remain ergodic at all investigated temperature and the temperature dependence of the corresponding characteristic time is well described by an Arrhenius law. Finally, we discuss the evidence in favor of the presence of a type-*A* transition temperature for the odd rotational degrees of freedom, distinct from the type-*B* transition associated with the arrest of the translational and even rotational ones, as predicted by the mode-coupling theory for the glass transition.

PACS numbers: 64.70.Pf, 61.20.Lc, 61.20.Ja - Version: February 2, 2008

The ideal mode-coupling theory (MCT) equations have been recently solved in the site-site representation for a system of symmetric hard dumbbell molecules [1, 2], as a function of the packing fraction φ and the elongation ζ . Interestingly enough, the theory predicts two different dynamic arrest scenarios, on varying φ and ζ (see Fig. 1 in Ref. [1]). For large elongations, it is predicted that all rotational correlation functions are strongly coupled to the translational degrees of freedom and dynamic arrest takes place at a common φ value, $\varphi_c^B(\zeta)$. According to MCT, the transition is of type-*B*, i.e. the long time limit of translational and rotational correlation functions jumps discontinuously from zero to a finite value at the ideal glass transition line. The ideal glass transition line (the *B*-line) has a non-monotonic shape in the φ - ζ plane, with a maximum at $\zeta = 0.43$. The *B*-line continues for small elongations until the hard sphere limit at $\zeta = 0$ is reached. However, for small elongations $\zeta < \zeta_c = 0.345$, theory predicts a novel different scenario: only the translational and the *even* rotational degrees of freedom freeze at the *B*-line. Here *even* and *odd* refer to the parity of the order l of the rotational correlator $C_l(t) = \langle P_l(\hat{e}(t) \cdot \hat{e}(0)) \rangle$, where P_l is the Legendre l -polynomial and $\hat{e}(t)$ is a unity vector along the molecular axis at time t . The *odd*- l rotational degrees of freedom freeze at higher values of the packing fraction, namely at an *A*-line $\varphi_c^A(\zeta) > \varphi_c^B(\zeta)$, that merges with the *B*-line at $\zeta = \zeta_c$. The glass transition of the odd degrees is of type-*A*, i.e. the long time limit of the odd rotational correlation functions increases continuously from zero on crossing the *A*-line.

The *A*- and *B*-lines separate the φ - ζ plane in three dynamic regions: an ergodic fluid, a completely arrested state, and an intermediate state (located between the *A*- and *B*-lines) which is the amorphous analog of a plastic crystal. In such a state, each molecule remains trapped in the cage formed by the neighbouring molecules, but

is able to perform 180° rotations within the cage, which lead to relaxation for the odd- l but not for the even- l rotational correlators. An analogous scenario is predicted by molecular MCT for the case of hard-ellipsoids in the limit of small aspect ratio [3].

In order to test these predictions, we have recently carried out molecular dynamics simulations [4] in a binary mixture of Lennard-Jones (LJ) dumbbell molecules at fixed packing fraction, as a function of the elongation using the temperature T as control parameter (instead of φ used in the theoretical work). In agreement with MCT predictions, a non-monotonic dependence on the elongation has been observed for the isodiffusivity curves, with a minimum at a value of ζ close to the maximum of $\varphi_c^B(\zeta)$ [1]. For small elongations $\zeta \leq 0.3$, in deep supercooled states, the coherent intermediate scattering function evaluated at the maximum of the static structure factor, as well as the rotational correlator $C_2(t)$, decays to zero at times several orders of magnitude longer than those for $C_1(t)$. An analysis of the plateau height for the $C_1(t)$ correlator shows a sharp drop of the former in the range $0.3 < \zeta < 0.5$, consistent with the theoretical ζ_c value predicted for the hard dumbbell fluid. These features suggest the presence of a nearby type-*A* transition in the investigated LJ dumbbell system.

Simulations in Ref. [4] have been carried out in *equilibrium* liquid states, for state points above the *B*-transition, i.e., at T such that translational, and both the odd and even rotational degrees of freedom are ergodic. In this manuscript we complement equilibrium simulations and we also investigate the reorientational dynamics *below* the *B*-transition, i.e., in *out-of-equilibrium* states, where the translational and even rotational degrees of freedom are arrested. We aim at providing evidence of the type-*A* transition, and to shed light on the dynamic features of amorphous analogs to plastic crystals.




FIG. 1: Isodiffusivity lines (thin continuous lines), and τ_1 (dotted lines) and τ_2 (thin dashed lines) isochrones in the T - ζ plane. These quantities are averaged over all particles in the system. Line segments join the numerical data (symbols are not included for clarity). The corresponding values (from top to bottom) are: $D = 1 \times 10^{-2}, 3 \times 10^{-3}, 1 \times 10^{-3}, 3 \times 10^{-4}, 1 \times 10^{-4}, 3 \times 10^{-5}$; $\tau_1 = 3 \times 10^1, 1 \times 10^2, 3 \times 10^2, 1 \times 10^3, 3 \times 10^3, 1 \times 10^4, 3 \times 10^4, 1 \times 10^5$; $\tau_2 = 3 \times 10^1, 1 \times 10^2, 3 \times 10^2, 1 \times 10^3, 3 \times 10^3, 1 \times 10^4$. Empty and filled symbols are respectively the estimated MCT and VFT temperatures (see text) for D (circles), τ_1 (squares), and τ_2 (triangles). Thick dashed and thick continuous lines are corresponding guides for the eyes.

We investigate a binary mixture of 410:102 dumbbell molecules [5] in a box of side L with periodic boundary conditions. Each molecule consists of two identical atoms, denoted by A and B respectively for the most and the less abundant component of the mixture. The interaction between atoms of different molecules is given by a Lennard-Jones potential plus a linear term: $V_{\alpha\beta}(r) = 4\epsilon_{\alpha\beta}[(\sigma_{\alpha\beta}/r)^{12} - (\sigma_{\alpha\beta}/r)^6 + A_0 + A_1(r/\sigma_{\alpha\beta})]$, where $\alpha, \beta \in (A, B)$. The interaction parameters are the same of the monoatomic binary mixture of Kob and Andersen [6], i.e., $\epsilon_{AA} = 1.0$, $\epsilon_{AB} = 1.5$, $\epsilon_{BB} = 0.5$, $\sigma_{AA} = 1.0$, $\sigma_{AB} = 0.8$, $\sigma_{BB} = 0.88$. The values of A_0 and A_1 are selected to guarantee continuity of potential and forces at the cutoff distance $c\sigma_{\alpha\beta}$, with $c = 2.5$. The resulting values are $A_0 = c^{-6}(7 - 13c^{-6})$ and $A_1 = 6c^{-7}(2c^{-6} - 1)$. The same atomic mass $m = 1$ is used for both A and B particles (each dumbbell has mass equal to two).

Simulations have been carried out for elongations $\zeta = 0.125, 0.20, 0.30, 0.40, 0.55, 0.70$ and 0.85 at different values of T and constant packing fraction $\varphi = 0.708$. The packing fraction is defined by $\varphi = (\pi/6)(N_{AA}\sigma_{AA}^3 + N_{BB}\sigma_{BB}^3)L^{-3}[1 + (3/2)\zeta - (1/2)\zeta^3]$, where $N_{AA} = 410$, $N_{BB} = 102$ are the number of AA and BB molecules. Temperature, time and diffusion coefficient D are respectively calculated in units of ϵ_{AA}/k_B , $(m\sigma_{AA}^2/\epsilon_{AA})^{1/2}$ and $(\sigma_{AA}^2\epsilon_{AA}/m)^{1/2}$. Equations of motion are integrated,

with a time step ranging from 2×10^{-4} to 5×10^{-3} according to T and/or ζ , using the velocity Verlet scheme combined with the SHAKE algorithm [7] for keeping the molecular elongation constant. At each state point, the system is thermalized at the requested T by periodic velocity rescaling. A microcanonical run is performed after suitable equilibration to generate a representative trajectory in phase space. Statistical averages are performed over 20 independently generated equilibrium trajectories. At high T , the thermalization process is started from a distorted crystalline configuration that quickly melts. At low T , it is started from equilibrated high T configurations. After reaching equilibrium, energy or pressure show no drift. Also, translational and rotational mean squared displacements as well as dynamic correlation functions, show no aging, i.e. there is no systematic dependence of the dynamical properties from the selected time origin. In practice equilibration requires that each dumbbell, in average, has moved several diameters and performed several full rotations from its initial position.

At very low T and small ζ , relaxation times for translational and even rotational degrees of freedom become exceedingly long for the time scale of the simulation. In these conditions only the odd rotational degrees of freedom decay to zero. We perform long runs of $\approx 10^7$ steps until neither significant drift in T and pressure, nor aging in the translational and even rotational correlation functions is observed, to ensure that the residual rotational motion of the dumbbells takes place in an essentially frozen structure [8]. After this preliminary simulation, trajectories are calculated.

Fig. 1 shows data for D and for the relaxation times τ_1 , τ_2 of the rotational correlators $C_1(t)$ and $C_2(t)$. The diffusion coefficient is evaluated from the long time limit of the mean square displacement, while τ_1 and τ_2 are defined as the time at which the corresponding correlation functions decay to 3 % of their initial value. As previously done in the analysis of other systems characterized by slow dynamics [4, 9], we report isochrones (constant τ_i) and isodiffusivity (constant D) curves, to provide an indication of the shape of the glass line in the $(T - \zeta)$ plane. The isodiffusivity curves are consistent with the same quantities calculated in Ref. [4] and, as previously noted, show a minimum around $\zeta = 0.55$, close to the MCT prediction for hard dumbbells [1]. Fig. 1 shows that, especially at low T , the iso- τ_2 curves follow the same trend as the isodiffusivity curves, tending to collapse in a common curve, as expected on approaching a type-B transition. The iso- τ_1 curves show a similar trend only at large ζ . On the contrary, for small elongations the iso- τ_1 curves depart from the iso- D and iso- τ_2 curves. Differently from τ_2 , within the resolution in ζ of our calculations, the τ_1 isochrones are monotonic in the $T - \zeta$ plane. The fact that the iso- τ_1 curves lay significantly below the iso- τ_2 curves shows very clearly that fast relaxation of the *odd* degrees of freedom takes place even when the even degrees are arrested.

A detailed quantitative comparison of the presented re-




FIG. 2: Symbols: Temperature dependence of D , τ_1 and τ_2 . Continuous lines are fits to VFT functions (see text). Dashed lines are Arrhenius fits. Data for τ_1 have been represented vs. $1/T$ in order to evidence the small ζ Arrhenius behavior.

sults with MCT predictions for dynamic exponents and scaling laws can not be simply performed, except for large ζ . Indeed, it has been shown that the range of validity of the asymptotic MCT laws — for the small ζ hard dumbbell fluid case [2] — is limited to unusually small values of the separation parameter $\epsilon = (\varphi_c^B - \varphi)/\varphi_c^B$. In particular, the exponent γ for the well-known power law $D, 1/\tau \propto (\varphi_c^B - \varphi)^\gamma$ can only be correctly estimated by exploring the range $\epsilon < 10^{-2}$. For small ζ , a fit of data in the range $10^{-2} < \epsilon < 10^{-1}$ would show an apparent power law behavior, but the corresponding fit would yield an “effective” exponent, which contrary to the asymptotic limit, would not take a universal value, but would depend on the chosen observable [2]. A similar phenomenon is expected to be present in the studied system at small ζ , preventing the possibility of an accurate determination of the asymptotic laws close to the MCT critical temperature T_c^B . In addition, we recall that ideal

MCT neglects activated processes which are known to be present in reality around and below T_c^B [10]. Only when hopping phenomena can be neglected, an MCT study can be feasible very close to T_c^B .

Fig. 2 shows the T dependence of D , τ_1 and τ_2 for several ζ values. If a fit (not shown) of the T dependence of D and $1/\tau_2$ according to the functional form $(T - T_c^B)^\gamma$ is performed, we obtain very similar values of T_c^B for both quantities in all the ζ range (see Fig. 1). For the case of $1/\tau_1$, reasonable power-law fits can be performed only for $\zeta \gtrsim 0.4$. For $\zeta = 0.55$ fits provide values of T_c^B consistent with those obtained for D and $1/\tau_2$ but with different γ values. Only for $\zeta \geq 0.70$ a simultaneous consistent description for D , $1/\tau_1$ and $1/\tau_2$ is achieved, with identical T_c^B and γ values. Specifically $T_c^B = 0.732$, $\gamma = 2.50$ for $\zeta = 0.70$ and $T_c^B = 1.02$, $\gamma = 2.06$ for $\zeta = 0.85$.

To provide a lower bound to T_c^B at small ζ , we also fit the data in terms of the well-known phenomenological Vogel-Fulcher-Tammann (VFT) law $D, 1/\tau_{1,2} \propto \exp[-A/(T - T_{VFT})]$, which predicts a dynamic arrest of the corresponding degree of freedom at a finite temperature $T_{VFT} < T_c^B$ [10]. In all cases, the VFT law provides a consistent effective representation of the dynamics, with a fit quality comparable to the MCT power law fit. The resulting $T_{VFT}(\zeta)$ for D , τ_1 and τ_2 are reported in Fig. 1. Interestingly, T_{VFT} for D , τ_1 and τ_2 coincide for large elongations, indicating strong couplings between the translational and all the rotational degrees of freedom. However, for $\zeta \lesssim 0.4$ the T_{VFT} for τ_1 is much smaller than the T_{VFT} for D and τ_2 , confirming that around this value of ζ the odd degrees of freedom completely decouple from the even and from the translational ones. This latter value of ζ is consistent with the merging point of the A - and B -transitions for the hard dumbbell fluid [1] and with estimates of Ref. [4]. At smaller elongations $\zeta \leq 0.3$, for τ_1 we find $T_{VFT} \approx 0$. This implies that a good description of the data is provided by an Arrhenius law $\tau_1 \propto \exp(E/k_B T)$ (see Fig. 2). The activation energies $E/\epsilon_{AA} = 1.8, 3.3$ and 5.0 , respectively for $\zeta = 0.125, 0.20$ and 0.30 , roughly follow a linear dependence on ζ . The presence of an activated Arrhenius T dependence in $\tau_1(T)$ for small molecular elongations, strongly suggests that the residual reorientational dynamics at very low temperatures, in the T -region where the translational dynamics is already arrested, is characterized by 180° rotations which do not lead to relaxation of the even rotational degrees of freedom. The Arrhenius functional form also suggests that hopping phenomena are extremely relevant, so much to completely mask the presence of a finite type- A glass transition temperature, if the analysis is limited only to the T dependence of τ_1 (or any other odd τ).

While we can not expect, due to this intense hopping decorrelation, to observe any critical law, indication of a residual cross-over finite temperature can still be observed in the full dependence of the correlation functions, and in particular in the strength of the long time α relax-




FIG. 3: Temperature dependence of the rotational correlators $C_2(t)$ and $C_3(t)$ vs. time rescaled by $T^{1/2}$, for elongation $\zeta = 0.2$. From top to bottom $T = 0.217, 0.276, 0.343, 0.496, 0.750, 1.110, 1.396, 1.693, 1.770, 1.815, 1.933, 2.443$ and 3.776 .

ation process. The main effect of hopping is indeed manifested in the long time decay from the plateau value. Indeed, in the absence of these additional relaxation channels, the correlation function would decay only down to the plateau value. Fig. 3 shows, for $\zeta = 0.2$, the T -dependence of $C_2(t)$ and $C_3(t)$, in a wide range of T , (across $T_{VTF} = 0.96$ for τ_2). Though $C_1(t)$ has more experimental relevance, $C_3(t)$ is more adequate to discuss the behavior of the non-ergodicity parameter, since it shows a more pronounced plateau. A representation of $C_i(t)$ vs. $tT^{1/2}$, i.e. rescaling the time by the thermal velocity, yields a clearer visualization of the decay of the correlation function. Data reported in Fig. 3 show two important features:

(i) at low T the plateau height (i.e., the non-ergodicity factor) for both C_2 and C_3 progressively grows up, signaling a crossing of the critical MCT temperatures. However, while in case of $C_2(t)$, the increase of the plateau

value is connected to a non-ergodic long time behavior (i.e. C_2 does not decay to zero within the simulation time window), $C_3(t)$ always relaxes toward zero. This fact evidences that hopping processes restore ergodicity for reorientations at T far below the temperature at which the plateau value starts to increase from zero. (ii) The decay of the C_2 correlation function is characterized by a finite T interval in which the plateau value is essentially constant (within the present numerical error). For smaller T , the height of the plateau starts to increase, as discussed above. Instead, the decay of the C_3 correlation function is not characterized by any plateau at high T , and the increase of the plateau starts from zero, consistently with the presence of a type-A transition which, due to the hopping effects, is significantly weakened. Moreover, the first T at which the C_3 plateau value is different from zero is lower than the T at which the C_2 plateau starts to increase. This suggests that the type-A transition temperature is different and lower than the type-B T_c^B . We also note that the α -relaxation process for odd C_i is highly stretched, which might reflect the local cage heterogeneity.

In summary, we have provided evidence for the existence of the type-A transition predicted by MCT for the reorientational dynamics of glass-forming liquids of small molecular elongation. Our numerical results complement those recently reported for $T > T_c^B$ in Ref. [4]. By computing relaxation times and plateau heights of odd rotational dynamic correlators — in the arrested state for the even rotational and the translational degrees of freedom — we show that reorientational hopping dominates the dynamics and restores ergodicity down to temperatures far below the type-A transition. Though these results have been specifically obtained for a liquid of symmetric dumbbells, experimental techniques probing $C_1(t)$ and $C_2(t)$, at very low temperatures, will yield qualitatively similar results for small elongated molecules with a weak asymmetry which disfavours rotations different from $\approx 180^\circ$. Finally, the present model can become a test case for developing theories of the glass transition in which activated processes — which become relevant around and below T_c^B — can be studied.

MIUR-FIRB is acknowledged for financial support.

-
- [1] S.-H. Chong and W. Götze, Phys. Rev. E **65**, 041503 (2002).
 - [2] S.-H. Chong and W. Götze, Phys. Rev. E **65**, 051201 (2002).
 - [3] M. Letz, R. Schilling, and A. Latz, Phys. Rev. E **62**, 5173 (2000).
 - [4] S.-H. Chong, A. J. Moreno, F. Sciortino, and W. Kob, Phys. Rev. Lett. (to be published).
 - [5] The system investigated in Ref. [4] is a 800:200 mixture with a modified LJ interaction.
 - [6] W. Kob and H. C. Andersen, Phys. Rev. Lett. **73**, 1376 (1994).
 - [7] M. P. Allen and D. J. Tildesley, *Computer Simulations*

- of Liquids*, Oxford University Press, Oxford, UK (1997).
- [8] In principle, significant aging effects for translational and even rotational degrees of freedom could be observed for much longer simulation times, beyond the limit of current computational resources.
- [9] E. Zaccarelli *et al.*, Phys. Rev. E **66**, 041402 (2002); G. Foffi *et al.*, Phys. Rev. Lett. **90**, 238301 (2003).
- [10] S. Kämmerer, W. Kob, and R. Schilling, Phys. Rev. E **56**, 5450 (1997); M. Aichele and J. Baschnagel, Eur. Phys. J. E **5**, 245 (2001); S. S. Ashwin and S. Sastry, J. Phys.: Condens. Matter **15**, S1253 (2003).